

What is Hyperbolic Geometry?

Mahan Mj,
School of Mathematics,
Tata Institute of Fundamental Research.

History

Euclid's Axioms

- 1 Any two points in a plane may be joined by a straight line.
- 2 A finite straight line may be extended continuously in a straight line.
- 3 A circle may be constructed with any centre and radius.
- 4 All right angles are equal to one another.
- 5 If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which the angles are less than the two right angles.

History

Euclid's Axioms

- 1 Any two points in a plane may be joined by a straight line.
- 2 A finite straight line may be extended continuously in a straight line.
- 3 A circle may be constructed with any centre and radius.
- 4 All right angles are equal to one another.
- 5 If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which the angles are less than the two right angles.

History

Euclid's Axioms

- 1 Any two points in a plane may be joined by a straight line.
- 2 A finite straight line may be extended continuously in a straight line.
- 3 A circle may be constructed with any centre and radius.
- 4 All right angles are equal to one another.
- 5 If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which the angles are less than the two right angles.

History

Euclid's Axioms

- 1 Any two points in a plane may be joined by a straight line.
- 2 A finite straight line may be extended continuously in a straight line.
- 3 A circle may be constructed with any centre and radius.
- 4 All right angles are equal to one another.
- 5 If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which the angles are less than the two right angles.

History

Euclid's Axioms

- 1 Any two points in a plane may be joined by a straight line.
- 2 A finite straight line may be extended continuously in a straight line.
- 3 A circle may be constructed with any centre and radius.
- 4 All right angles are equal to one another.
- 5 If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which the angles are less than the two right angles.

History

Euclid's Axioms

- 1 Any two points in a plane may be joined by a straight line.
- 2 A finite straight line may be extended continuously in a straight line.
- 3 A circle may be constructed with any centre and radius.
- 4 All right angles are equal to one another.
- 5 If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which the angles are less than the two right angles.

Circa 100 BC:

5th postulate is equivalent to

Through a point not on a straight line there is one and only one straight line through the point parallel to the given straight line.

The attempt to prove the 5th postulate from the other postulates gave rise to hyperbolic geometry.

Circa 100 BC:

5th postulate is equivalent to

Through a point not on a straight line there is one and only one straight line through the point parallel to the given straight line.

The attempt to prove the 5th postulate from the other postulates gave rise to hyperbolic geometry.

Circa 100 BC:

5th postulate is equivalent to

Through a point not on a straight line there is one and only one straight line through the point parallel to the given straight line.

The attempt to prove the 5th postulate from the other postulates gave rise to hyperbolic geometry.

Circa 100 BC:

5th postulate is equivalent to

Through a point not on a straight line there is one and only one straight line through the point parallel to the given straight line.

The attempt to prove the 5th postulate from the other postulates gave rise to hyperbolic geometry.

History

Gauss started thinking of parallels about 1792. In an 18th November, 1824 letter to F. A. Taurinus, he wrote:

‘The assumption that the sum of the three angles (of a triangle) is smaller than two right angles leads to a geometry which is quite different from our (Euclidean) geometry, but which is in itself completely consistent.

But Gauss did not publish his work.

History

Gauss started thinking of parallels about 1792. In an 18th November, 1824 letter to F. A. Taurinus, he wrote:

'The assumption that the sum of the three angles (of a triangle) is smaller than two right angles leads to a geometry which is quite different from our (Euclidean) geometry, but which is in itself completely consistent.

But Gauss did not publish his work.

History

Gauss started thinking of parallels about 1792. In an 18th November, 1824 letter to F. A. Taurinus, he wrote:

‘The assumption that the sum of the three angles (of a triangle) is smaller than two right angles leads to a geometry which is quite different from our (Euclidean) geometry, but which is in itself completely consistent.

But Gauss did not publish his work.

History

Gauss started thinking of parallels about 1792. In an 18th November, 1824 letter to F. A. Taurinus, he wrote:

‘The assumption that the sum of the three angles (of a triangle) is smaller than two right angles leads to a geometry which is quite different from our (Euclidean) geometry, but which is in itself completely consistent.

But Gauss did not publish his work.

History

Gauss started thinking of parallels about 1792. In an 18th November, 1824 letter to F. A. Taurinus, he wrote:

‘The assumption that the sum of the three angles (of a triangle) is smaller than two right angles leads to a geometry which is quite different from our (Euclidean) geometry, but which is in itself completely consistent.

But Gauss did not publish his work.

History

Gauss started thinking of parallels about 1792. In an 18th November, 1824 letter to F. A. Taurinus, he wrote:

‘The assumption that the sum of the three angles (of a triangle) is smaller than two right angles leads to a geometry which is quite different from our (Euclidean) geometry, but which is in itself completely consistent.

But Gauss did not publish his work.

In the 18th century, Johann Heinrich Lambert introduced what are today called hyperbolic functions and computed the area of a hyperbolic triangle.

In the nineteenth century, hyperbolic geometry was extensively explored by the Hungarian mathematician Janos Bolyai and the Russian mathematician Nikolai Ivanovich Lobachevsky, after whom it is sometimes named. Lobachevsky published a paper entitled *On the principles of geometry* in 1829-30, while Bolyai discovered hyperbolic geometry and published his independent account of non-Euclidean geometry in the paper *The absolute science of space* in 1832.

The term "hyperbolic geometry" was introduced by Felix Klein in 1871.

In the 18th century, Johann Heinrich Lambert introduced what are today called hyperbolic functions and computed the area of a hyperbolic triangle.

In the nineteenth century, hyperbolic geometry was extensively explored by the Hungarian mathematician Janos Bolyai and the Russian mathematician Nikolai Ivanovich Lobachevsky, after whom it is sometimes named. Lobachevsky published a paper entitled *On the principles of geometry* in 1829-30, while Bolyai discovered hyperbolic geometry and published his independent account of non-Euclidean geometry in the paper *The absolute science of space* in 1832.

The term "hyperbolic geometry" was introduced by Felix Klein in 1871.

In the 18th century, Johann Heinrich Lambert introduced what are today called hyperbolic functions and computed the area of a hyperbolic triangle.

In the nineteenth century, hyperbolic geometry was extensively explored by the Hungarian mathematician Janos Bolyai and the Russian mathematician Nikolai Ivanovich Lobachevsky, after whom it is sometimes named. Lobachevsky published a paper entitled *On the principles of geometry* in 1829-30, while Bolyai discovered hyperbolic geometry and published his independent account of non-Euclidean geometry in the paper *The absolute science of space* in 1832.

The term "hyperbolic geometry" was introduced by Felix Klein in 1871.

In the 18th century, Johann Heinrich Lambert introduced what are today called hyperbolic functions and computed the area of a hyperbolic triangle.

In the nineteenth century, hyperbolic geometry was extensively explored by the Hungarian mathematician Janos Bolyai and the Russian mathematician Nikolai Ivanovich Lobachevsky, after whom it is sometimes named. Lobachevsky published a paper entitled *On the principles of geometry* in 1829-30, while Bolyai discovered hyperbolic geometry and published his independent account of non-Euclidean geometry in the paper *The absolute science of space* in 1832.

The term "hyperbolic geometry" was introduced by Felix Klein in 1871.

The Problem

Parallel Postulate

Given a straight line L in a plane P and a point x on the plane P lying outside the line L , there exists a unique straight line L' lying on P passing through x and parallel to L .

Problem

Prove the Parallel Postulate from the other axioms of Euclidean geometry.

The Problem

Parallel Postulate

Given a straight line L in a plane P and a point x on the plane P lying outside the line L , there exists a unique straight line L' lying on P passing through x and parallel to L .

Problem

Prove the Parallel Postulate from the other axioms of Euclidean geometry.

The Problem

Parallel Postulate

Given a straight line L in a plane P and a point x on the plane P lying outside the line L , there exists a unique straight line L' lying on P passing through x and parallel to L .

Problem

Prove the Parallel Postulate from the other axioms of Euclidean geometry.

The Problem

Parallel Postulate

Given a straight line L in a plane P and a point x on the plane P lying outside the line L , there exists a unique straight line L' lying on P passing through x and parallel to L .

Problem

Prove the Parallel Postulate from the other axioms of Euclidean geometry.

The Problem

Parallel Postulate

Given a straight line L in a plane P and a point x on the plane P lying outside the line L , there exists a unique straight line L' lying on P passing through x and parallel to L .

Problem

Prove the Parallel Postulate from the other axioms of Euclidean geometry.

The Problem

Parallel Postulate

Given a straight line L in a plane P and a point x on the plane P lying outside the line L , there exists a unique straight line L' lying on P passing through x and parallel to L .

Problem

Prove the Parallel Postulate from the other axioms of Euclidean geometry.

The Problem

Parallel Postulate

Given a straight line L in a plane P and a point x on the plane P lying outside the line L , there exists a unique straight line L' lying on P passing through x and parallel to L .

Problem

Prove the Parallel Postulate from the other axioms of Euclidean geometry.

The Problem

Parallel Postulate

Given a straight line L in a plane P and a point x on the plane P lying outside the line L , there exists a unique straight line L' lying on P passing through x and parallel to L .

Problem

Prove the Parallel Postulate from the other axioms of Euclidean geometry.

Question

What is:

- 1) a plane P*
- 2) a straight line L*
- 3) the notion of parallelism?*

Question

What is:

- 1) a plane P*
- 2) a straight line L*
- 3) the notion of parallelism?*

Question

What is:

- 1) a plane P*
- 2) a straight line L*
- 3) the notion of parallelism?*

Question

What is:

- 1) a plane P*
- 2) a straight line L*
- 3) the notion of parallelism?*

Question

What is:

- 1) a plane P*
- 2) a straight line L*
- 3) the notion of parallelism?*

Question

What is:

- 1) a plane P*
- 2) a straight line L*
- 3) the notion of parallelism?*

Euclidean Geometry Revisited

Answers for Euclidean Geometry:

The Euclidean plane is \mathbb{R}^2 equipped with the metric

$$ds^2 = dx^2 + dy^2.$$

(Infinitesimal Pythagoras)

Meaning: Lengths of curves σ (= *smooth* maps of $[0, 1]$ into \mathbb{R}^2) are computed as per the formula

$$l(\sigma) = \int_0^1 ds = \int_0^1 \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right]^{\frac{1}{2}} dt \quad (A)$$

for some parametrization $x = x(t)$, $y = y(t)$ of the curve σ .

Euclidean Geometry Revisited

Answers for Euclidean Geometry:

The Euclidean plane is \mathbb{R}^2 equipped with the **metric**

$$ds^2 = dx^2 + dy^2.$$

(Infinitesimal Pythagoras)

Meaning: Lengths of curves σ (= *smooth* maps of $[0, 1]$ into \mathbb{R}^2) are computed as per the formula

$$l(\sigma) = \int_0^1 ds = \int_0^1 \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right]^{\frac{1}{2}} dt \quad (A)$$

for some parametrization $x = x(t), y = y(t)$ of the curve σ .

Euclidean Geometry Revisited

Answers for Euclidean Geometry:

The Euclidean plane is \mathbb{R}^2 equipped with the **metric**

$$ds^2 = dx^2 + dy^2.$$

(Infinitesimal Pythagoras)

Meaning: Lengths of curves σ (= *smooth* maps of $[0, 1]$ into \mathbb{R}^2) are computed as per the formula

$$l(\sigma) = \int_0^1 ds = \int_0^1 \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right]^{\frac{1}{2}} dt \quad (A)$$

for some parametrization $x = x(t), y = y(t)$ of the curve σ .

Euclidean Geometry Revisited

Answers for Euclidean Geometry:

The Euclidean plane is \mathbb{R}^2 equipped with the **metric**

$$ds^2 = dx^2 + dy^2.$$

(Infinitesimal Pythagoras)

Meaning: Lengths of curves σ (= *smooth* maps of $[0, 1]$ into \mathbb{R}^2) are computed as per the formula

$$l(\sigma) = \int_0^1 ds = \int_0^1 \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right]^{\frac{1}{2}} dt \quad (A)$$

for some parametrization $x = x(t), y = y(t)$ of the curve σ .

Euclidean Geometry Revisited

Answers for Euclidean Geometry:

The Euclidean plane is \mathbb{R}^2 equipped with the **metric**

$$ds^2 = dx^2 + dy^2.$$

(Infinitesimal Pythagoras)

Meaning: Lengths of curves σ (= *smooth* maps of $[0, 1]$ into \mathbb{R}^2) are computed as per the formula

$$l(\sigma) = \int_0^1 ds = \int_0^1 \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right]^{\frac{1}{2}} dt \quad (A)$$

for some parametrization $x = x(t), y = y(t)$ of the curve σ .

Euclidean Geometry Revisited

Answers for Euclidean Geometry:

The Euclidean plane is \mathbb{R}^2 equipped with the **metric**

$$ds^2 = dx^2 + dy^2.$$

(Infinitesimal Pythagoras)

Meaning: Lengths of curves σ (= *smooth* maps of $[0, 1]$ into \mathbb{R}^2) are computed as per the formula

$$l(\sigma) = \int_0^1 ds = \int_0^1 \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right]^{\frac{1}{2}} dt \quad (A)$$

for some parametrization $x = x(t), y = y(t)$ of the curve σ .

Euclidean Geometry Revisited

Answers for Euclidean Geometry:

The Euclidean plane is \mathbb{R}^2 equipped with the **metric**

$$ds^2 = dx^2 + dy^2.$$

(Infinitesimal Pythagoras)

Meaning: Lengths of curves σ (= *smooth* maps of $[0, 1]$ into \mathbb{R}^2) are computed as per the formula

$$l(\sigma) = \int_0^1 ds = \int_0^1 \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right]^{\frac{1}{2}} dt \quad (A)$$

for some parametrization $x = x(t), y = y(t)$ of the curve σ .

Euclidean Geometry Revisited

Answers for Euclidean Geometry:

The Euclidean plane is \mathbb{R}^2 equipped with the **metric**

$$ds^2 = dx^2 + dy^2.$$

(Infinitesimal Pythagoras)

Meaning: Lengths of curves σ (= *smooth* maps of $[0, 1]$ into \mathbb{R}^2) are computed as per the formula

$$l(\sigma) = \int_0^1 ds = \int_0^1 \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right]^{\frac{1}{2}} dt \quad (A)$$

for some parametrization $x = x(t), y = y(t)$ of the curve σ .

Euclidean Geometry Revisited

Answers for Euclidean Geometry:

The Euclidean plane is \mathbb{R}^2 equipped with the **metric**

$$ds^2 = dx^2 + dy^2.$$

(Infinitesimal Pythagoras)

Meaning: Lengths of curves σ (= *smooth* maps of $[0, 1]$ into \mathbb{R}^2) are computed as per the formula

$$l(\sigma) = \int_0^1 ds = \int_0^1 \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right]^{\frac{1}{2}} dt \quad (A)$$

for some parametrization $x = x(t)$, $y = y(t)$ of the curve σ .

Euclidean Geometry Revisited

Answers for Euclidean Geometry:

The Euclidean plane is \mathbb{R}^2 equipped with the **metric**

$$ds^2 = dx^2 + dy^2.$$

(Infinitesimal Pythagoras)

Meaning: Lengths of curves σ (= *smooth* maps of $[0, 1]$ into \mathbb{R}^2) are computed as per the formula

$$l(\sigma) = \int_0^1 ds = \int_0^1 \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right]^{\frac{1}{2}} dt \quad (A)$$

for some parametrization $x = x(t)$, $y = y(t)$ of the curve σ .

Theorem

Given two points $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, the straight line segment between (x_1, y_1) and (x_2, y_2) is the unique path that realizes the shortest distance (as per formula A) between them.

Definition

Two bi-infinite straight lines are said to be parallel if they do not intersect.

Theorem

Given two points $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, the straight line segment between (x_1, y_1) and (x_2, y_2) is the unique path that realizes the shortest distance (as per formula A) between them.

Definition

Two bi-infinite straight lines are said to be parallel if they do not intersect.

Theorem

Given two points $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, the straight line segment between (x_1, y_1) and (x_2, y_2) is the unique path that realizes the shortest distance (as per formula A) between them.

Definition

Two bi-infinite straight lines are said to be parallel if they do not intersect.

General Geometries

Answers for a slightly more general geometry: Metric on $U \subset \mathbb{R}^2$:

$$ds^2 = f(x, y)dx^2 + g(x, y)dy^2.$$

Here, $l(\sigma) = \int_0^1 ds =$

$$\int_0^1 [f(x(t), y(t))\left(\frac{dx}{dt}\right)^2 + g(x(t), y(t))\left(\frac{dy}{dt}\right)^2]^{\frac{1}{2}} dt \dots (B)$$

General Geometries

Answers for a slightly more general geometry: Metric on $U \subset \mathbb{R}^2$:

$$ds^2 = f(x, y)dx^2 + g(x, y)dy^2.$$

Here, $l(\sigma) = \int_0^1 ds =$

$$\int_0^1 [f(x(t), y(t))\left(\frac{dx}{dt}\right)^2 + g(x(t), y(t))\left(\frac{dy}{dt}\right)^2]^{\frac{1}{2}} dt \dots (B)$$

General Geometries

Answers for a slightly more general geometry: Metric on $U \subset \mathbb{R}^2$:

$$ds^2 = f(x, y)dx^2 + g(x, y)dy^2.$$

Here, $l(\sigma) = \int_0^1 ds =$

$$\int_0^1 [f(x(t), y(t))\left(\frac{dx}{dt}\right)^2 + g(x(t), y(t))\left(\frac{dy}{dt}\right)^2]^{\frac{1}{2}} dt \dots (B)$$

General Geometries

Answers for a slightly more general geometry: Metric on $U \subset \mathbb{R}^2$:

$$ds^2 = f(x, y)dx^2 + g(x, y)dy^2.$$

Here, $l(\sigma) = \int_0^1 ds =$

$$\int_0^1 [f(x(t), y(t))\left(\frac{dx}{dt}\right)^2 + g(x(t), y(t))\left(\frac{dy}{dt}\right)^2]^{\frac{1}{2}} dt \dots (B)$$

General Geometries

Answers for a slightly more general geometry: Metric on $U \subset \mathbb{R}^2$:

$$ds^2 = f(x, y)dx^2 + g(x, y)dy^2.$$

Here, $l(\sigma) = \int_0^1 ds =$

$$\int_0^1 [f(x(t), y(t))\left(\frac{dx}{dt}\right)^2 + g(x(t), y(t))\left(\frac{dy}{dt}\right)^2]^{\frac{1}{2}} dt \dots (B)$$

General Geometries

Answers for a slightly more general geometry: Metric on $U \subset \mathbb{R}^2$:

$$ds^2 = f(x, y)dx^2 + g(x, y)dy^2.$$

Here, $l(\sigma) = \int_0^1 ds =$

$$\int_0^1 [f(x(t), y(t))\left(\frac{dx}{dt}\right)^2 + g(x(t), y(t))\left(\frac{dy}{dt}\right)^2]^{\frac{1}{2}} dt \dots (B)$$

Definitions:

- 1) Given two points $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, the **geodesic** between (x_1, y_1) and (x_2, y_2) is the unique path that realizes the shortest distance (as per formula B) between them.
- 2) An **isometry** I is a map that preserves the metric, i.e. if $I((x, y)) = (x_1, y_1)$ then

$$f(x, y)dx^2 + g(x, y)dy^2 = f(x_1, y_1)dx_1^2 + g(x_1, y_1)dy_1^2.$$

Definitions:

- 1) Given two points $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, the **geodesic** between (x_1, y_1) and (x_2, y_2) is the unique path that realizes the shortest distance (as per formula B) between them.
- 2) An **isometry** I is a map that preserves the metric, i.e. if $I((x, y)) = (x_1, y_1)$ then

$$f(x, y)dx^2 + g(x, y)dy^2 = f(x_1, y_1)dx_1^2 + g(x_1, y_1)dy_1^2.$$

Definitions:

- 1) Given two points $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, the **geodesic** between (x_1, y_1) and (x_2, y_2) is the unique path that realizes the shortest distance (as per formula B) between them.
- 2) An **isometry** I is a map that preserves the metric, i.e. if $I((x, y)) = (x_1, y_1)$ then

$$f(x, y)dx^2 + g(x, y)dy^2 = f(x_1, y_1)dx_1^2 + g(x_1, y_1)dy_1^2.$$

Definitions:

1) Given two points $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, the **geodesic** between (x_1, y_1) and (x_2, y_2) is the unique path that realizes the shortest distance (as per formula B) between them.

2) An **isometry** I is a map that preserves the metric, i.e. if $I((x, y)) = (x_1, y_1)$ then

$$f(x, y)dx^2 + g(x, y)dy^2 = f(x_1, y_1)dx_1^2 + g(x_1, y_1)dy_1^2.$$

Definitions:

- 1) Given two points $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, the **geodesic** between (x_1, y_1) and (x_2, y_2) is the unique path that realizes the shortest distance (as per formula B) between them.
- 2) An **isometry** I is a map that preserves the metric, i.e. if $I((x, y)) = (x_1, y_1)$ then

$$f(x, y)dx^2 + g(x, y)dy^2 = f(x_1, y_1)dx_1^2 + g(x_1, y_1)dy_1^2.$$

Definitions:

- 1) Given two points $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, the **geodesic** between (x_1, y_1) and (x_2, y_2) is the unique path that realizes the shortest distance (as per formula B) between them.
- 2) An **isometry** I is a map that preserves the metric, i.e. if $I((x, y)) = (x_1, y_1)$ then

$$f(x, y)dx^2 + g(x, y)dy^2 = f(x_1, y_1)dx_1^2 + g(x_1, y_1)dy_1^2.$$

Definitions:

- 1) Given two points $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, the **geodesic** between (x_1, y_1) and (x_2, y_2) is the unique path that realizes the shortest distance (as per formula B) between them.
- 2) An **isometry** I is a map that preserves the metric, i.e. if $I((x, y)) = (x_1, y_1)$ then

$$f(x, y)dx^2 + g(x, y)dy^2 = f(x_1, y_1)dx_1^2 + g(x_1, y_1)dy_1^2.$$

Hyperbolic Geometry

Answers for hyperbolic geometry:

A **Model** for hyperbolic geometry is the upper half plane

$\mathbf{H} = (x, y) \in \mathbb{R}^2, y > 0$ equipped with the metric
 $ds^2 = \frac{1}{y^2}(dx^2 + dy^2)$.

Where does this come from? Another story.

Hyperbolic Geometry

Answers for hyperbolic geometry:

A Model for hyperbolic geometry is the upper half plane

$\mathbf{H} = (x, y) \in \mathbb{R}^2, y > 0$ equipped with the metric
 $ds^2 = \frac{1}{y^2}(dx^2 + dy^2)$.

Where does this come from? Another story.

Hyperbolic Geometry

Answers for hyperbolic geometry:

A Model for hyperbolic geometry is the upper half plane

$\mathbf{H} = (x, y) \in \mathbb{R}^2, y > 0$ equipped with the metric

$$ds^2 = \frac{1}{y^2}(dx^2 + dy^2).$$

Where does this come from? Another story.

Hyperbolic Geometry

Answers for hyperbolic geometry:

A Model for hyperbolic geometry is the upper half plane

$\mathbf{H} = (x, y) \in \mathbb{R}^2, y > 0$ equipped with the metric
 $ds^2 = \frac{1}{y^2}(dx^2 + dy^2)$.

Where does this come from? Another story.

Hyperbolic Geometry

Answers for hyperbolic geometry:

A Model for hyperbolic geometry is the upper half plane

$\mathbf{H} = (x, y) \in \mathbb{R}^2, y > 0$ equipped with the metric
 $ds^2 = \frac{1}{y^2}(dx^2 + dy^2)$.

Where does this come from? *Another story.*

Hyperbolic Geometry

Answers for hyperbolic geometry:

A Model for hyperbolic geometry is the upper half plane

$\mathbf{H} = (x, y) \in \mathbb{R}^2, y > 0$ equipped with the metric
 $ds^2 = \frac{1}{y^2}(dx^2 + dy^2)$.

Where does this come from? Another story.

Hyperbolic Geometry

Answers for hyperbolic geometry:

A Model for hyperbolic geometry is the upper half plane

$\mathbf{H} = (x, y) \in \mathbb{R}^2, y > 0$ equipped with the metric
 $ds^2 = \frac{1}{y^2}(dx^2 + dy^2)$.

Where does this come from? Another story.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z .

Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic.
Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic.
Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .

Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .

Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.

Theorem 1:

Vertical straight lines in \mathbf{H} are geodesics. In fact, the vertical segment between a, b is the **unique** geodesic between a, b .


Theorem 2:

1) *Translations*: Define $f : \mathbf{H} \rightarrow \mathbf{H}$ by $f(x, y) = (x + a, y)$ for some fixed $a \in \mathbb{R}$.

2) *Inversions about semicircles*: Define $g : \mathbf{H} \rightarrow \mathbf{H}$ by $g(z) = \frac{R^2}{\bar{z}}$ for some $R > 0$, where \bar{z} denotes the complex conjugate of z . Then f, g are isometries of \mathbf{H} .


Observation 3:

Image of a geodesic under an isometry is another geodesic. Hence images of vertical geodesics under inversions are geodesics.



Inversions about semicircles

$$I(\overline{ab}) \cdot I(\overline{ac}) = I(\overline{ad})^2.$$



Inversions about semicircles

$$I(\overline{ab}) \cdot I(\overline{ac}) = I(\overline{ad})^2.$$

Theorem 4: In the hyperbolic plane \mathbf{H} ,
vertical straight lines
and semi-circular arcs with center on the real axis
are geodesics.

Theorem 4: In the hyperbolic plane \mathbf{H} ,
vertical straight lines
and semi-circular arcs with center on the real axis
are geodesics.

Theorem 4: In the hyperbolic plane \mathbf{H} ,
vertical straight lines
and semi-circular arcs with center on the real axis
are geodesics.

Theorem 4: In the hyperbolic plane \mathbf{H} ,
vertical straight lines
and semi-circular arcs with center on the real axis
are geodesics.

Theorem 4: In the hyperbolic plane \mathbf{H} ,
vertical straight lines
and semi-circular arcs with center on the real axis
are geodesics.

Theorem 4: In the hyperbolic plane \mathbf{H} ,
vertical straight lines
and semi-circular arcs with center on the real axis
are geodesics.

We have the following **New Parallel Postulate**:

Given a geodesic L in \mathbf{H} and a point x on \mathbf{H} lying outside L , there exist **INFINITELY** many bi-infinite geodesics L' lying on \mathbf{H} passing through x and parallel to L .

We have the following **New Parallel Postulate**:

Given a geodesic L in \mathbf{H} and a point x on \mathbf{H} lying outside L , there exist **INFINITELY** many bi-infinite geodesics L' lying on \mathbf{H} passing through x and parallel to L .

We have the following **New Parallel Postulate**:

Given a geodesic L in \mathbf{H} and a point x on \mathbf{H} lying outside L , there exist **INFINITELY** many bi-infinite geodesics L' lying on \mathbf{H} passing through x and parallel to L .

We have the following **New Parallel Postulate**:

Given a geodesic L in \mathbf{H} and a point x on \mathbf{H} lying outside L , there exist **INFINITELY** many bi-infinite geodesics L' lying on \mathbf{H} passing through x and parallel to L .

We have the following **New Parallel Postulate**:

Given a geodesic L in \mathbf{H} and a point x on \mathbf{H} lying outside L , there exist **INFINITELY** many bi-infinite geodesics L' lying on \mathbf{H} passing through x and parallel to L .

We have the following **New Parallel Postulate**:

Given a geodesic L in \mathbf{H} and a point x on \mathbf{H} lying outside L , there exist **INFINITELY** many bi-infinite geodesics L' lying on \mathbf{H} passing through x and parallel to L .

We have the following **New Parallel Postulate**:

Given a geodesic L in \mathbf{H} and a point x on \mathbf{H} lying outside L , there exist **INFINITELY** many bi-infinite geodesics L' lying on \mathbf{H} passing through x and parallel to L .